PATENT APPLICATION

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of

Masashi OGAWA et al.

Continuation of Appln. No.: 09/125,944

Group Art Unit in Parent: 1631

Confirmation No.: Not Yet Assigned

Examiner in Parent: M. MORAN

Filed: July 31, 2001

For:

METHOD OF MEASUREMENT OF PROTEASE AND THIN MEMBRANES USED

FOR SAID METHOD

PRELIMINARY AMENDMENT

Commissioner for Patents Washington, D.C. 20231

Sir:

Prior to examination, please amend the above-identified application as follows:

IN THE SPECIFICATION:

Please amend the specification by inserting before the first line the sentence:

--This is a continuation of Application No. 09/125,944, filed February 10, 1999, the disclosure of which is incorporated herein by reference.--

IN THE CLAIMS:

Please enter the following amended claims:

- 1. (amended) A method for measuring protease which comprises the steps of:
- (1) contacting a sample containing protease with a thin membrane which comprises a protease substrate together with a hardening agent formed on a surface of a support; and

- (2) detecting a trace of digestion formed on the thin membrane by the action of protease.
 - 2. (amended) A method for measuring protease which comprises the steps of:
- (1) contacting one of two substantially continuous slices of a biological sample with a thin membrane which comprises a protease substrate together with a hardening agent formed on a surface of a support;
- (2) detecting a trace of digestion formed on the thin membrane by the action of protease; and
- (3) comparing the trace of digestion with a histopathological preparation prepared from the other slice.
 - 3. (amended) A method for measuring protease which comprises the steps of:
- (1) contacting one of two or more substantially continuous slices of a biological sample with a thin membrane which comprises a protease substrate together with a hardening agent formed on a surface of a support:
- (2) contacting the remaining slices with a thin membrane which comprises a protease substrate, a hardening agent, and a protease inhibitor formed on a surface of a support;
- (3) detecting traces of digestion formed on the thin membranes by the action of protease; and
- (4) comparing the trace of digestion on the thin membrane used in step (1) with the trace of digestion on the thin membrane used in step (2).
 - 4. (amended) A method for measuring protease which comprises the steps of:

- (1) contacting one of two or more substantially continuous slices of a biological sample with a thin membrane which comprises a protease substrate together with a hardening agent formed on a surface of a support;
- (2) contacting the remaining slices with a thin membrane which comprises a protease substrate different from the protease substrate present in the thin membrane used in step (1) together with a hardening agent formed on a surface of a support;
- (3) detecting traces of digestion formed on the thin membranes by the action of protease; and
- (4) comparing the trace of digestion on the thin membrane used in step (1) with the trace of digestion on the thin membrane used in step (2).
 - 5. (amended) A method for measuring protease which comprises the steps of:
- (1) contacting a sample containing protease with a thin membrane which comprises at least the following two layers: layer (a) which contains a protease substrate, a hardening agent, and a protease inhibitor formed on a surface of a support, and layer (b) which contains a protease substrate and a hardening agent laminated on layer (a);
- (2) detecting traces of digestion formed on the thin membrane by the action of protease; and
- (3) comparing the trace of digestion on layer (a) with the trace of digestion on layer (b).
 - 6. (amended) A method for measuring protease which comprises the steps of:
- (1) contacting a sample containing protease with a thin membrane which comprises at least the following two layers: layer (a) which contains a protease substrate together with a

hardening agent formed on a surface of a support, and layer (b) which contains a protease substrate different from the protease substrate present in layer (a) together with a hardening agent laminated on layer (a);

- (2) detecting traces of digestion formed on the thin membrane by the action of protease; and
- (3) comparing the trace of digestion on layer (a) with the trace of digestion on layer (b).
- 7. (amended) The method of <u>claim 1</u> wherein the protease substrate is selected from the group consisting of collagen, gelatin, proteoglycan, fibronectin, laminin, elastin, and casein.
- 8. (amended) The method of <u>claim 1</u> wherein the sample is a biological sample isolated or collected from a patient.
- 9. (amended) The method of <u>claim 1</u> wherein the detecting by using a thin membrane containing one or more substances selected from the group consisting of metals, metal oxides, pigments and dyes and having a maximum transmission density of 0.01 or higher at a wavelength ranging from 400 nm to 700 nm.
- 10. (amended) The method of claim 1 wherein the protease is a matrix metalloproteinase.
- 11. (amended) A thin membrane for measuring protease which contains a protease substrate together with a hardening agent formed on a surface of a support.
- 12. (amended) The thin membrane of claim 11 which comprises at least the following two layers:

layer (a) which comprises a protease substrate, a hardening agent and a protease inhibitor formed on a surface of a support, and layer (b) which contains a protease substrate together with a hardening agent laminated on layer (a).

- 13. (amended) The thin membrane of claim 11 which comprises at least the following two layers: layer (a) which comprises a protease substrate together with a hardening agent formed on a surface of a support, and layer (b) which comprises a protease substrate different from the protease substrate present in layer (a) together with a hardening agent laminated on layer (a).
- 14. (amended) The thin membrane of claim 11 which comprises one or more substances selected from the group consisting of metals, metal oxides, pigments and dyes and have a maximum transmission density of 0.01 or higher at a wavelength ranging from 400 nm to 700 nm.
- 15. (amended) The thin membrane of <u>claim 11</u> wherein the support is a microscope slide or a polyethylene terephthalate film.
- 16. (amended) The thin membrane of <u>claim 11</u> wherein an undercoat layer is present between the support and the thin membrane.
- 17. (amended) A method of diagnosing a disease involving protease which comprises the steps of:
- (1) contacting a biological sample isolated or collected from a patient with a thin membrane which comprises a protease substrate together with a hardening agent formed on a surface of a support; and

- (2) detecting the trace of digestion formed on the thin membrane by the action of protease.
- 18. (amended) The method of claim 17 wherein the disease is selected from the group consisting of cancer, rheumatic diseases, periodontal diseases and alveolar pyorrhea.

REMARKS

Claims 1-18 have been amended to remove improper multiple dependencies and for editorial purposes only. No new matter has been added.

Entry and consideration of this Amendment is respectfully requested.

Respectfully submitted,

Drew Hissong

Registration No. 44,765

SUGHRUE, MION, ZINN, MACPEAK & SEAS, PLLC 2100 Pennsylvania Avenue, N.W. Washington, D.C. 20037-3213 Telephone: (202) 293-7060 Facsimile: (202) 293-7860

Date: July 31, 2001

APPENDIX

VERSION WITH MARKINGS TO SHOW CHANGES MADE

IN THE CLAIMS:

The claims are amended as follows:

- 1. (amended) A method for measuring protease which comprises the steps of:
- (1) <u>contacting bringing</u> a sample containing protease into contact with a thin membrane which comprises a protease substrate together with a hardening agent and is formed on a surface of a support; and
- (2) detecting the <u>a</u> trace of digestion formed on the thin membrane by the action of protease.
 - 2. (amended) A method for measuring protease which comprises the steps of:
- (1) <u>contacting bringing</u> one of two substantially continuous slices of a biological sample into contact with a thin membrane which comprises a protease substrate together with a hardening agent and is-formed on a surface of a support;
- (2) detecting the <u>a</u> trace of digestion formed on the thin membrane by the action of protease; and
- (3) comparing the trace of digestion with a histopathological preparation prepared from the other slice.
 - 3. (amended) A method for measuring protease which comprises the steps of:
- (1) <u>contacting bringing</u> one of two or more substantially continuous slices of a biological sample into contact with a thin membrane which comprises a protease substrate together with a hardening agent and is formed on a surface of a support;

- (2) <u>contacting bringing</u> the remaining slices into contact with a thin membrane which comprises a protease substrate, a hardening agent, and a protease inhibitor and is-formed on a surface of a support;
- (3) detecting traces of digestion formed on the thin membranes by the action of protease; and
- (4) comparing the trace of digestion on the thin membrane used in the step (1) with the trace of digestion on the thin membrane used in the step (2).
 - 4. (amended) A method for measuring protease which comprises the steps of:
- (1) <u>contacting bringing</u> one of two or more substantially continuous slices of a biological sample into contact with a thin membrane which comprises a protease substrate together with a hardening agent and is formed on a surface of a support;
- (2) <u>contacting bringing</u> the remaining slices into contact with a thin membrane which comprises a protease substrate different from the protease substrate contained <u>present</u> in the thin membrane used in the step (1) together with a hardening agent and is formed on a surface of a support;
- (3) detecting traces of digestion formed on the thin membranes by the action of protease; and
- (4) comparing the trace of digestion on the thin membrane used in the step (1) with the trace of digestion on the thin membrane used in the step (2).
 - 5. (amended) A method for measuring protease which comprises the steps of:
- (1) <u>contacting bringing</u> a sample containing protease into contact with a thin membrane which comprises at least the following two layers: layer (a) which contains a protease

PRELIMINARY AMENDMENT

Continuation of U.S. Appln. No. 09/125,944

substrate, a hardening agent, and a protease inhibitor and is-formed on a surface of a support, and layer (b) which contains a protease substrate and a hardening agent and is-laminated on the-layer (a);

- (2) detecting traces of digestion formed on the thin membrane by the action of protease; and
- (3) comparing the trace of digestion on the layer (a) with the trace of digestion on the layer (b).
 - 6. (amended) A method for measuring protease which comprises the steps of:
- (1) <u>contacting bringing</u> a sample containing protease <u>into contact</u> with a thin membrane which comprises at least the following two layers: layer (a) which contains a protease substrate together with a hardening agent and is formed on a surface of a support, and layer (b) which contains a protease substrate different from the protease substrate <u>present contained</u> in the layer (a) together with a hardening agent and is laminated on the layer (a);
- (2) detecting traces of digestion formed on the thin membrane by the action of protease; and
- (3) comparing the trace of digestion on the layer (a) with the trace of digestion on the layer (b).
- 7. (amended) The method of <u>claim 1</u> any one of claims 1 to 6 wherein the protease substrate is selected from the group consisting of collagen, gelatin, proteoglycan, fibronectin, laminin, elastin, and casein.
- 8. (amended) The method of <u>claim 1</u> any one of claims 1 to 7 wherein the sample is a biological sample isolated or collected from a patient.

- 9. (amended) The method of <u>claim 1</u> any one of claims 1 to 8 wherein the <u>detecting</u> detection is performed by using a thin membrane containing one or more substances selected from the group consisting of metals, metal oxides, pigments and dyes and having a maximum transmission density of 0.01 or higher at a wavelength ranging from 400 nm to 700 nm.
- 10. (amended) The method of <u>claim 1 any one of claims 1 to 9</u> wherein the protease is <u>a</u> matrix metalloproteinase.
- 11. (amended) A thin membrane for measuring protease which contains a protease substrate together with a hardening agent and is formed on a surface of a support.
- 12. (amended) The thin membrane of claim 11 which comprises at least the following two layers:

layer (a) which comprises a protease substrate, a hardening agent and a protease inhibitor and is-formed on a surface of a support, and layer (b) which contains a protease substrate together with a hardening agent and is-laminated on the-layer (a).

- 13. (amended) The thin membrane of claim 11 which comprises at least the following two layers: layer (a) which comprises a protease substrate together with a hardening agent and is formed on a surface of a support, and layer (b) which comprises a protease substrate different from the protease substrate contained present in the layer (a) together with a hardening agent and is laminated on the layer (a).
- 14. (amended) The thin membrane of <u>claim 11</u> any one of <u>claims 11 to 13</u> which <u>comprise comprises</u> one or more substances selected from the group consisting of metals, metal oxides, pigments and dyes and have a maximum transmission density of 0.01 or higher at a wavelength ranging from 400 nm to 700 nm.

- 15. (amended) The thin membrane of <u>claim 11 any one of claims 11 to 14</u> wherein the support is <u>selected from</u> a microscope slide <u>and or a polyethylene terephthalate film</u>.
- 16. (amended) The thin membrane of <u>claim 11 any one of claims 11 to 15</u> wherein an undercoat layer is present provided between the support and the thin membrane.
- 17. (amended) A method of diagnosing a disease involving protease which comprises the steps of:
- (1) <u>contacting bringing</u> a biological sample isolated or collected from a patient into eontact with a thin membrane which comprises a protease substrate together with a hardening agent and is-formed on a surface of a support; and
- (2) detecting the trace of digestion formed on the thin membrane by the action of protease.
- 18. (amended) The method of <u>claim 17</u> wherein the disease is selected from the group consisting of cancer, rheumatic diseases, periodontal diseases and alveolar pyorrhea.